

MATHEMATICS


Chapter 4: PRINCIPLE OF MATHEMATICAL INDUCTION




PRINCIPLE OF MATHEMATICAL INDUCTION

Top Concepts

1. There are two types of reasoning—**deductive** and **inductive**.
2. In deduction, given a statement to be proven which is often called a conjecture or a theorem, valid deductive steps are derived and a proof may or may not be established.
3. Deduction is the application of a general case to a particular case.
4. Inductive reasoning depends on working with each case and developing a conjecture by observing incidence till each and every case is observed.
5. Induction is the generalisation from particular cases or facts.
6. A deductive approach is known as a 'top-down approach'. Given the theorem which is narrowed down to specific *hypotheses* then to *observation*. Finally, the hypotheses is tested with specific data to get the *confirmation* (or not) of original theory.



7. Inductive reasoning works the other way—moving from specific observations to broader generalisations and theories. Informally, this is known as a 'bottom-up approach'.




8. To prove statements or results formulated in terms of n , where n is a positive integer, a principle based on inductive reasoning called the **Principle of Mathematical Induction (PMI)** is used.
9. PMI is one such tool which can be used to prove a wide variety of mathematical statements. Each of such statements is assumed as $P(n)$ associated with a positive integer n for which the correctness of the case $n = 1$ is examined. Then, assuming the truth of $P(k)$ for some positive integer k , the truth of $P(k + 1)$ is established.
10. Let $p(n)$ denote a mathematical statement such that
 - (1) $p(1)$ is true.
 - (2) $p(k + 1)$ is true whenever $p(k)$ is true.
 Then, the statement is true for all natural numbers n by PMI.
11. PMI is based on the Peano's Axiom.
12. PMI is based on a series of well-defined steps, so it is necessary to verify all of them.
13. PMI can be used to prove the equality, inequalities and divisibility of natural numbers.

Key Formulae

1. Sum of n natural numbers: $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$
2. Sum of n^2 natural numbers: $1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$
3. Sum of odd natural numbers: $1 + 3 + 5 + 7 + \dots + (2n - 1) = n^2$
4. Steps of PMI
 1. Denote the given statement in terms of n by $P(n)$.
 2. Check whether the proposition is true for $n = 1$.
 3. Assume that the proposition result is true for $n = k$.
 4. Using $p(k)$, prove that the proposition is true for $p(k + 1)$.
5. Rules of inequalities
 - a. If $a < b$ and $b < c$, then $a < c$.
 - b. If $a < b$, then $a + c < b + c$.
 - c. If $a < b$ and $c > 0$ which means c is positive, then $ac < bc$.
 - d. If $a < b$ and $c < 0$ which means c is negative, then $ac > bc$.

MIND MAP : LEARNING MADE SIMPLE CHAPTER - 4



Important Questions

Multiple Choice questions-

Question 1. For all $n \in \mathbb{N}$, $3n^5 + 5n^3 + 7n$ is divisible by

- (a) 5
- (b) 15
- (c) 10
- (d) 3

Question 2. $\{1 - (1/2)\}\{1 - (1/3)\}\{1 - (1/4)\} \dots \{1 - 1/(n + 1)\} =$

- (a) $1/(n + 1)$ for all $n \in \mathbb{N}$.
- (b) $1/(n + 1)$ for all $n \in \mathbb{R}$
- (c) $n/(n + 1)$ for all $n \in \mathbb{N}$.
- (d) $n/(n + 1)$ for all $n \in \mathbb{R}$

Question 3. For all $n \in \mathbb{N}$, $3^{2n} + 7$ is divisible by

- (a) non of these
- (b) 3
- (c) 11
- (d) 8

Question 4. The sum of the series $1 + 2 + 3 + 4 + 5 + \dots n$ is

- (a) $n(n + 1)$
- (b) $(n + 1)/2$
- (c) $n/2$
- (d) $n(n + 1)/2$

Question 5. The sum of the series $1^2 + 2^2 + 3^2 + \dots n^2$ is

- (a) $n(n + 1) (2n + 1)$
- (b) $n(n + 1) (2n + 1)/2$
- (c) $n(n + 1) (2n + 1)/3$
- (d) $n(n + 1) (2n + 1)/6$

Question 6. For all positive integers n , the number $n(n^2 - 1)$ is divisible by:

- (a) 36

(b) 24

(c) 6

(d) 16

Question 7. If n is an odd positive integer, then $a^n + b^n$ is divisible by :

(a) $a^2 + b^2$

(b) $a + b$

(c) $a - b$

(d) none of these

Question 8. $n(n + 1)(n + 5)$ is a multiple of _____ for all $n \in \mathbb{N}$

(a) 2

(b) 3

(c) 5

(d) 7

Question 9. For any natural number n , $7^n - 2^n$ is divisible by

(a) 3

(b) 4

(c) 5

(d) 7

Question 10. The sum of the series $1^3 + 2^3 + 3^3 + \dots + n^3$ is

(a) $\{(n + 1)/2\}^2$

(b) $\{n/2\}^2$

(c) $n(n + 1)/2$

(d) $\{n(n + 1)/2\}^2$

Very Short:

1.

Short Questions:

1. For every integer n , prove that $7n - 3n$ divisible by 4.

2. Prove that $n(n + 1)(n + 5)$ is multiple of 3.

3. Prove that $10^{2n-1} + 1$ is divisible by 11.

4. Prove that $\left(1 + \frac{1}{1}\right)\left(1 + \frac{1}{2}\right)\left(1 + \frac{1}{3}\right) \dots \left(1 + \frac{1}{n}\right) = (n + 1)$

5. Prove $1.2 + 2.3 + 3.4 + \dots + n(n+1) = \frac{n(n+1)(n+2)}{3}$

Long Questions:

1. Prove $(2n+7) < (n+3)^2$

2. Prove that:

$$\frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3n-2)(3n+1)} = \frac{n}{(3n+1)}$$

3. Prove $1.2 + 2.22 + 3.23 + \dots + n.2^n = (n-1)^{2n+1} + 2$

4. Prove that $2.7^n + 3.5^n - 5$ is divisible by 24 $\forall n \in \mathbb{N}$.

5. Prove that $41^n - 14^n$ is a multiple of 27.

Answer Key:

MCQ:

1. (b) 15

2. (a) $1/(n+1)$ for all $n \in \mathbb{N}$.

3. (d) 8

4. (d) $n(n+1)/2$

5. (d) $n(n+1)(2n+1)/6$

6. (c) 6

7. (b) $a+b$

8. (b) 3

9. (c) 5

10. (d) $\{n(n+1)/2\}^2$

Very Short Answer:

1. $\left(\frac{\pi}{32}\right)^c$

2. $39^\circ 22' 30''$

3. $\frac{5\pi}{12} \text{ cm}$

4. $\sqrt{3}$

5. $\frac{-1}{\sqrt{2}}$

6. $2 - \sqrt{3}$

7. $\frac{-4}{5}$

8. 45°

9. $2 \sin 8\theta \cos 4\theta$

10. $\sin 6x - \sin 2x$

Short Answer:

1. $P(n) : 7^n - 3^n$ is divisible by 4

For $n = 1$

$P(1) : 7^1 - 3^1 = 4$ which is divisible by 4. Thus, $P(1)$ is true

Let $P(k)$ be true

$7^k - 3^k$ is divisible by 4

$7^k - 3^k = 4\lambda$, where $\lambda \in \mathbb{N}$ (i)

we want to prove that $P(k+1)$ is true whenever $P(k)$ is true

$$7^{k+1} - 3^{k+1} = 7^k \cdot 7 - 3^k \cdot 3$$

$$= (4\lambda + 3^k) \cdot 7 - 3^k \cdot 3 \text{ (from i)}$$

$$= 28\lambda + 7 \cdot 3^k - 3^k \cdot 3$$

$$= 28\lambda + 3^k(7 - 3)$$

$$= 4(7\lambda + 3^k)$$

Hence

$7^{k+1} - 3^{k+1}$ is divisible by 4

thus $P(k+1)$ is true when $P(k)$ is true.

Therefore by P.M.I. the statement is true for every positive integer n .

- 2.

$P(n) : n(n+1)(n+5)$ is multiple of 3

for $n=1$

$P(1) : 1(1+1)(1+5) = 12$ is multiple of 3

let $P(k)$ be true

$P(k) : k(k+1)(k+5)$ is multiple of 3

$$\Rightarrow k(k+1)(k+5) = 3\lambda \text{ where } \lambda \in \mathbb{N} \text{ (i)}$$

we want to prove that result is true for $n=k+1$

$P(k+1) : (k+1)(k+2)(k+6)$

$$\begin{aligned}
 &\Rightarrow (K+1)(k+2)(k+6) = [(k+1)(k+2)](k+6) \\
 &= k(k+1)(k+2) + 6(k+1)(k+2) \\
 &= k(k+1)(k+5-3) + 6(k+1)(k+2) \\
 &= k(k+1)(k+5) - 3k(k+1) + 6(k+1)(K+2) \\
 &= k(k+1)(k+5) + (k+1)[6(k+2) - 3k] \\
 &= k(k+1)(k+5) + (k+1)(3k+12) \\
 &= k(k+1)(k+5) + 3(k+1)(k+4) \\
 &= 3\lambda + 3(k+1)(k+4) \text{ (from i)} \\
 &= 3[\lambda + (K+1)(K+4)] \text{ which is multiple of three} \\
 &\text{Hence } P(k+1) \text{ is multiple of 3.}
 \end{aligned}$$

3.

$$P(n): 10^{2n-1} + 1 \text{ is divisible by 11}$$

$$\text{for } n=1$$

$$P(1) = 10^{2 \times 1 - 1} + 1 = 11 \text{ is divisible by 11. Hence result is true for } n=1$$

let $P(k)$ be true

$$P(k): 10^{2k-1} + 1 \text{ is divisible by 11}$$

$$\Rightarrow 10^{2k-1} + 1 = 11\lambda \text{ where } \lambda \in \mathbb{N}(i)$$

we want to prove that result is true for $n = k + 1$

$$= 10^{2(k+1)-1} + 1 = 10^{2k+2-1} + 1$$

$$= 10^{2k+1} + 1$$

$$= 10^{2k} \cdot 10^1 + 1$$

$$= (110\lambda - 10) \cdot 10 + 1 \text{ (from i)}$$

$$= 1100\lambda - 100 + 1$$

$$= 1100\lambda - 99$$

$$= 11(100\lambda - 9) \text{ is divisible by 11}$$

Hence by P.M.I. $P(k+1)$ is true whenever $P(k)$ is true.

4.

$$\text{let } P(n): \left(1 + \frac{1}{1}\right)\left(1 + \frac{1}{2}\right)\left(1 + \frac{1}{3}\right) \dots \left(1 + \frac{1}{n}\right) = (n+1)$$

$$\text{for } n=1$$

$$P(1): \left(1 + \frac{1}{1}\right) = (1+1) = 2$$

which is true

let $P(k)$ be true

$$P(k) : \left(1 + \frac{1}{1}\right)\left(1 + \frac{1}{2}\right)\left(1 + \frac{1}{3}\right) \dots \left(1 + \frac{1}{k}\right) = (k+1)$$

we want to prove that $P(k+1)$ is true

$$P(k+1) : \left(1 + \frac{1}{1}\right)\left(1 + \frac{1}{2}\right) \dots \left(1 + \frac{1}{k+1}\right) = (k+2)$$

$$L.H.S. = \left(1 + \frac{1}{1}\right)\left(1 + \frac{1}{2}\right) \dots \left(1 + \frac{1}{k}\right)\left(1 + \frac{1}{k+1}\right)$$

$$= (k+1)\left(1 + \frac{1}{k+1}\right) \quad [from(1)]$$

$$= (k+1)\left(\frac{k+1+1}{k+1}\right)$$

$$= (k+2)$$

thus $P(k+1)$ is true whenever

$P(k)$ is true.

5.

$$p(n) : 1.2 + 2.3 + \dots + n(n+1) = \frac{n(n+1)(n+2)}{3}$$

for $n=1$

$$p(1) : 1(1+1) = \frac{1(1+1)(1+2)}{3}$$

$$p(1) = 2 = 2$$

hence $p(1)$ be true

$$p(k) : 1.2 + 2.3 + \dots + k(k+1) = \frac{k(k+1)(k+2)}{3} \dots\dots\dots(i)$$

we want to prove that

$$p(k+1) :$$

$$1.2 + 2.3 + \dots + (k+1)(k+2) = \frac{(k+1)(k+2)(k+3)}{3}$$

$L.H.S.$

$$= 1.2 + 2.3 + \dots + k(k+1) + (k+1)(k+2)$$

$$= \frac{k(k+1)(k+2)}{3} + \frac{(k+1)(k+2)}{1} \quad [from(i)]$$

$$\frac{k(k+1)(k+2) + 3(k+1)(k+2)}{3}$$

$$\frac{(k+1)(k+2)[k+3]}{3}$$

hence $p(k+1)$ is true whenever $p(k)$ is true

Long Answer:

1.

$$p(n) : (2n+7) < (n+3)^2$$

for $n=1$

$$9 < (4)^2$$

$$9 < 16$$

which is true

let $p(k)$ be true

$$(2k+7) < (k+3)^2$$

now

$$2(k+1)+7 = (2k+7)+2$$

$$< (k+3)^2 + 2 = k^2 + 6k + 11$$

$$< k^2 + 8k + 16 = (k+4)^2$$

$$= (k+3+1)^2$$

$$\therefore p(k+1) : 2(k+1)+7 < (k+1+3)^2$$

$$\Rightarrow p(k+1) \text{ is true, whenever } p(k) \text{ is true}$$

hence by PMI $p(k)$ is true for all $n \in N$

2.

$$p(n) : \frac{1}{1.4} + \frac{1}{4.7} + \dots + \frac{1}{(3n-2)(3n+1)} = \frac{n}{(3n+1)}$$

for $n=1$

$$p(1) : \frac{1}{(3-2)(3+1)} = \frac{1}{(3+1)} = \frac{1}{4}$$

which is true

let $p(k)$ be true

$$p(k) : \frac{1}{1.4} + \frac{1}{4.7} + \dots + \frac{1}{(3k-2)(3k+1)} = \frac{k}{(3k+1)} \dots\dots\dots (i)$$

we want to prove that $p(k+1)$ is true

$$p(k+1): \frac{1}{1.4} + \frac{1}{4.7} + \dots + \frac{1}{(3k+1)(3k+4)} = \frac{k+1}{(3k+4)}$$

L.H.S.

$$= \frac{1}{1.4} + \frac{1}{4.7} + \dots + \frac{1}{(3k-2)(3k+1)} + \frac{1}{(3k+1)(3k+4)}$$

$$= \frac{k}{3k+1} + \frac{1}{(3k+1)(3k+4)} \quad [\text{from} \dots (i)]$$

$$= \frac{k(3k+4) + 1}{(3k+1)(3k+4)}$$

$$= \frac{3k^2 + 4k + 1}{(3k+1)(3k+4)} = \frac{\cancel{(3k+1)}(k+1)}{\cancel{(3k+1)}(3k+4)}$$

$p(k+1)$ is true whenever $p(k)$ is true.

3.

$$p(n): 1.2 + 2.2^2 + 3.2^3 + \dots + n.2^n = (n-1)2^{n+1} + 2$$

$$p(n): 1.2 + 2.2^2 + 3.2^3 + \dots + n.2^n = (n-1)2^{n+1} + 2$$

for $n=1$

$$p(1): 1.2^1 = (1-1)2^2 + 2$$

$$2 = 2 \text{ which is true}$$

let $p(k)$ be true

$$p(k): 1.2 + 2.2^2 + \dots + k.2^k = (k-1)2^{k+1} + 2 \dots (i)$$

we want to prove that $p(k+1)$ is true

$$p(k+1): 1.2 + 2.2^2 + \dots + (k+1)2^{k+1} = k.2^{k+2} + 2$$

L.H.S.

$$1.2 + 2.2^2 + \dots + k.2^k + (k+1)2^{k+1} \quad [\text{from} \dots (i)]$$

$$= (k-1)2^{k+1} + 2 + (k+1)2^{k+1}$$

$$= 2^{k+1}(k-1 + k+1) + 2$$

$$= 2^{k+2}k + 2$$

This $p(k+1)$ is true whenever $p(k)$ is true

4. $P(n): 2.7^n + 3.5^n - 5$ is divisible by 24

for $n=1$

$$P(1): 2.7^1 + 3.5^1 - 5 = 24 \text{ is divisible by 24}$$

Hence result is true for $n=1$

Let $P(K)$ be true

$$P(K) : 2 \cdot 7^K + 3 \cdot 5^K - 5$$

$$\Rightarrow 2 \cdot 7^K + 3 \cdot 5^K - 5 = 24\lambda \text{ when } \lambda \in \mathbb{N}$$

we want to prove that $P(K+1)$ is True whenever $P(K)$ is true

$$2 \cdot 7^{K+1} + 3 \cdot 5^{K+1} - 5 = 2 \cdot 7^K \cdot 7 + 3 \cdot 5^K \cdot 5 - 5$$

$$= 7[2 \cdot 7^K + 3 \cdot 5^K - 5 - 3 \cdot 5^K + 5] + 3 \cdot 5^K \cdot 5 - 5$$

$$= 7[24\lambda - 3 \cdot 5^K + 5] + 15 \cdot 5^K - 5 \text{ (from i)}$$

$$= 7 \times 24\lambda - 21 \cdot 5^K + 35 + 15 \cdot 5^K - 5$$

$$= 7 \times 24\lambda - 6 \cdot 5^K + 30$$

$$= 7 \times 24\lambda - 6(5^K - 5)$$

$$= 7 \times 24\lambda - 6 \cdot 4p \left[\because 5^K - 5 \text{ is multiple of } 4 \right]$$

$$= 24(7\lambda - p), \quad 24 \text{ is divisible by } 24$$

Hence by P M I $p(n)$ is true for all $n \in \mathbb{N}$.

5. $P(n) : 41^n - 14^n$ is a multiple of 27

for $n = 1$

$$P(1) : 41^1 - 14 = 27, \text{ which is a multiple of } 27$$

Let $P(K)$ be True

$$P(K) : 41^K - 14^K$$

$$\Rightarrow 41^K - 14^K = 27\lambda, \text{ where } \lambda \in \mathbb{N}$$

we want to prove that result is true for $n = K + 1$

$$41^{K+1} - 14^{K+1} = 41^K \cdot 41 - 14^K \cdot 14$$

$$= (27\lambda + 14^K) \cdot 41 - 14^K \cdot 14 \text{ (from i)}$$

$$= 27\lambda \cdot 41 + 14^K \cdot 41 - 14^K \cdot 14$$

$$= 27\lambda \cdot 41 + 14^K (41 - 14)$$

$$= 27\lambda \cdot 41 + 14^K (27)$$

$$= 27(41\lambda + 14^K) \quad \text{is a multiple of } 27$$

Hence by PMI $p(n)$ is true for all $n \in \mathbb{N}$.